



- SISTERS OF CHARITY HOSPITAL • Buffalo, NY SISTERS OF CHARITY HOSPITAL ST JOSEPH Campus • Cheektowaga, NY
- KENMORE MERCY HOSPITAL • Kenmore, NY MERCY HOSPITAL • Buffalo, NY MERCY HOSPITAL Orchard Park division • Orchard Park, NY

Allergies & Sensitivities:

- No Known

Ambulatory Surgery Antibiotics Orders

Authorization is hereby given to dispense the generic/therapeutic equivalent unless otherwise indicated by the prescriber

Date:

Time:

Prescriber Orders

PRE-OPERATIVE ORDERS

SURGICAL INFECTION PROPHYLAXIS

- Preoperative antibiotics are not indicated (rationale): _____
- The following antibiotic(s) is to be initiated within 60 minutes *prior* incision (120 minutes prior to incision for levofloxacin (Levaquin) and Vancomycin). Antibiotics should be discontinued within 24 hours postoperatively (for prophylaxis). See selection in matrix below.

Antibiotic	Recommended Prophylaxis
<input type="checkbox"/> Cefazolin (Ancef) 1 gram IV <input type="checkbox"/> Cefazolin (Ancef) 2 grams IV Penicillin/Cephalosporin allergy OR History of MRSA: <input type="checkbox"/> Vancomycin 1 gram IV	Surgical cases to include: Cardiac (Pacemakers/AICD), Neurosurgical, Orthopedic, Podiatry, Vascular
<input type="checkbox"/> Levofloxacin (Levaquin) 500 mg IV OR <input type="checkbox"/> Gentamicin IV 2 mg/kg [round to the nearest 10 mg] + Metronidazole (Flagyl) 500 mg IV	Surgical cases to include: Transrectal prostate biopsy
<input type="checkbox"/> Ampicillin/sulbactam (Unasyn) IV 1.5 gram OR Penicillin/Cephalosporin allergy OR History of MRSA: <input type="checkbox"/> Vancomycin 1 gram IV + gentamicin 2 mg/kg IV [round to the nearest 10 mg]	Surgical cases to include: Penile prosthesis revision, insertion, removal
<input type="checkbox"/> Cefazolin (Ancef) 1 gram IV Penicillin/Cephalosporin allergy OR History of MRSA: <input type="checkbox"/> Vancomycin 1 gram IV	Surgical cases to include: Removal epididymis lesion
<input type="checkbox"/> Cefazolin (Ancef) 1 gram IV <input type="checkbox"/> Cefazolin (Ancef) 2 grams IV Penicillin/Cephalosporin allergy OR History of MRSA: <input type="checkbox"/> Vancomycin 1 gram IV + gentamicin 2 mg/kg IV [round to the nearest 10 mg]	Surgical cases to include: Gastric/Biliary – Endoscopy PEG placement
<input type="checkbox"/> Cefazolin (Ancef) 1 gram IV <input type="checkbox"/> Cefoxitin (Mefoxin) 2 grams IV Penicillin/Cephalosporin allergy <input type="checkbox"/> Metronidazole (Flagyl) 500 mg IV + Gentamicin IV 2 mg/kg [round to the nearest 10 mg]	Surgical cases to include: Laparoscopic-assisted hysterectomy Vaginal hysterectomy Pubovaginal sling
<input type="checkbox"/> Cefazolin (Ancef) 1 gram IV OR <input type="checkbox"/> Clindamycin (Cleocin) 600 mg IV	Surgical cases to include: Head and Neck

Prescriber Signature: _____